

Evaluation of a Three-Dimensional Chemical Transport Model (PMCAMx) in the Mexico City Metropolitan Area

Spyros N. Pandis¹, Alexandra P. Tsibidi¹, Vlassis A. Karydis¹, Miguel Zavala², Wenfang Lei², Luisa T. Molina²

¹*Dept. of Chemical Engineering, University of Patras, Patras, Greece*

²*Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology (MIT), and Molina Center for Energy and the Environment (MCE2), USA.*

Atmospheric aerosols have adverse effects on human health, contribute to the visibility reduction and influence the energy balance of the planet. A three-dimensional chemical transport model (PMCAMx) (Gaydos et al., 2007) is used to simulate the particular matter (PM) mass composition distribution in the Mexico City Metropolitan Area (MCMA). PMCAMx uses the framework of CAMx (ENVIRON, 2002) modelling the processes of horizontal and vertical advection, horizontal and vertical dispersion, wet and dry deposition, and gas-phase chemistry. In addition to the above, PMCAMx includes three detailed aerosol modules: inorganic aerosol growth (Gaydos et al., 2003; Koo et al., 2003a), aqueous-phase chemistry (Fahey and Pandis, 2001), and secondary organic aerosol formation and growth (Koo et al., 2004). The aerosol thermodynamic model ISORROPIA has been improved as it now simulates explicitly the chemistry of Ca, Mg, and K salts and is linked to PMCAMx. The hybrid approach (Koo et al., 2003b) for modelling aerosol dynamics is applied in order to accurately simulate the inorganic components in coarse mode. This approach assumes that the smallest particles are in equilibrium while the condensation/evaporation equation is solved for the larger ones. The new CMU organic aerosol model, which is based on the splitting of the organic aerosol volatility range in discrete bins, is also used. The model predictions are evaluated against the PM and vapour concentration measurements from the MCMA-2003 Campaign (Molina et al., 2007).

References

Gaydos, T., Pinder, R., Koo, B., Fahey, K., Yarwood, G., and Pandis, S. N., (2007). Development and application of a three-dimensional Chemical Transport Model, PMCAMx. *Atmospheric Environment*, in press.

ENVIRON (2002). User's guide to the comprehensive air quality model with extensions (CAMx). Version 3.10. Report prepared by ENVIRON International corporation, Novato, CA

Gaydos, T., Koo, B., and Pandis, S. N., (2003). Development and application of an efficient moving sectional approach for the solution of the atmospheric aerosol condensation/evaporation equations. *Atmospheric Environment*, 37, 3303-3316.

Fahey, K. and Pandis, S. N., (2001). Optimizing model performance: variable size resolution in cloud chemistry modelling. *Atmospheric Environment* 35, 4471-4478.

Koo, B., Pandis S. N., and Ansari, A. (2003a). Integrated approaches to modelling the organic and inorganic atmospheric aerosol components. *Atmospheric Environment*, 37, 4757-4768.

Koo, B., Gaydos, T.M., Pandis, S.N., (2003b). Evaluation of the equilibrium, hybrid, and dynamic aerosol modeling approaches. *Aerosol Science and Technology* 37, 53–64

Molina, L.T., Kolb, C.E., de Foy, B., Lamb, B., Brune, W., Molina, M.J. Air Quality in North Americas Most Populous City Overview of MCMA-2003 Campaign. *Atmos. Chem. Phys. Discuss.* 7, 2007.